

BI-TP 97/07
 DESY 97-038
 WUE-ITP-97-02
 hep-ph/9703288

Leptoquark/Squark Interpretation of HERA Events: Virtual Effects in e^+e^- Annihilation to Hadrons

J. Kalinowski^{1,2}, R. Rückl^{3,*}, H. Spiesberger^{4,*},
 and P.M. Zerwas¹

¹ *Deutsches Elektronen-Synchrotron DESY, D-22607 Hamburg*

² *Institute of Theoretical Physics, Warsaw University, PL-00681 Warsaw*

³ *Institut für Theoretische Physik, Universität Würzburg, D-97074 Würzburg*

⁴ *Fakultät für Physik, Universität Bielefeld, D-33501 Bielefeld*

ABSTRACT

In reference to the recently observed high Q^2 , large x events in deep-inelastic positron–proton scattering at HERA, various leptoquark and supersymmetric scenarios are discussed. We study the impact of virtual leptoquark or R -parity breaking squark exchange as well as generic contact interaction on the production of quark–antiquark pairs in e^+e^- annihilation, in particular at LEP2.

* Supported by Bundesministerium für Bildung, Wissenschaft, Forschung und Technologie, Bonn, Germany, Contracts 05 7BI92P (9) and 05 7WZ91P (0).

1 Introduction

The recent observation of events in deep-inelastic positron-proton scattering with very high Q^2 and large x at HERA [1] has refuelled speculations on physics beyond the Standard Model, in particular on low-mass leptoquark-type particles. Such particles had been suggested a long time ago in a variety of physical scenarios: Pati-Salam $SU(4)$ unification of quarks and leptons [2], grand unified theories such as $SU(5)$ or E_6 [3], and composite models [4]. Moreover, in supersymmetric theories squarks couple to lepton-quark pairs if the R -symmetry is broken in the trilinear couplings of the superfields [5, 6]. Vector leptoquarks in grand unified theories with both lepton-quark and diquark couplings must be very heavy to suppress proton decay; certain scalar leptoquarks in GUT multiplets could nevertheless be relatively light [7] (disregarding the notorious hierarchy problem for the time being). Squarks in supersymmetric theories should naturally be expected in the mass range of a few hundred GeV.

A general classification of these novel states¹ has been presented in Ref. [8]. In this analysis the couplings of leptoquarks to lepton-quark pairs are assumed to be baryon- and lepton-number conserving in order to avoid rapid proton decay, family diagonal to exclude FCNC processes beyond the CKM mixing, and chiral to preserve the helicity suppression in leptonic pion decay. Moreover, the couplings are taken dimensionless and all interactions are assumed to respect the $SU(3)_C \times SU(2)_L \times U(1)_Y$ symmetry of the Standard Model. The allowed states can be classified according to spin, weak isospin and fermion number. We adopt the notation of Refs. [9] to conform with the notation generally employed in experimental papers: vector leptoquarks are denoted by V_I , scalar leptoquarks by S_I ; isomultiplets with different hypercharges are distinguished by a tilde.

For convenience the nine possible states of scalar and vector leptoquarks are listed in Table 1. Leptoquarks in the upper (lower) part of Table 1 carry fermion number $F = 2$ ($F = 0$). The couplings are denoted generically by g_R or g_L with R, L referring to the chirality of the lepton. Each state can couple with different strength; for simplicity, the additional indices are suppressed. In principle the two scalar states S_0 and $S_{1/2}$ and the two vector states V_0 and $V_{1/2}$ could have both chiral g_R and g_L couplings at the same time; however, since the product of the two couplings is constrained very strongly by rare decays [10, 11, 12, 13], we assume only one of the two couplings to be non-zero. The special type of leptoquark that does not induce proton decay (as a result of its quantum numbers) and is compatible with the renormalization of the electroweak mixing angle from the symmetry value $3/8$ at the GUT scale down to $\sin^2 \Theta_W \cong 0.23$ at the electroweak scale [7], is marked by an asterisk. Only a small subset of all these states is realized in supersymmetric theories with R -parity breaking. Moreover, supersymmetry requires these states to have universal left-handed couplings to leptons.

If leptoquarks or squarks in R -parity breaking supersymmetric theories exist, a large

¹We shall generically denote leptoquarks and squarks in R -parity breaking scenarios by LQ .

LQ		Q	Decay Mode	BR $e^\pm j$	Coupling	Limits Ref. [11, 12]	HERA estimates	$e_i^- e^+ \rightarrow q_k \bar{q}$ ik
S_0	$\tilde{d}_R$	$-1/3$	$e_L u$	$\frac{1}{2}$	g_L	$g_L < 0.06$	0.40	LL
			$\nu_L d$		$-g_L$			—
			$e_R u$	1	g_R	$g_R < 0.1$	0.28	RR
$\tilde{S}_0$		$-4/3$	$e_R d$	1	g_R	$g_R < 0.1$	0.30	RR
S_1		$+2/3$	$\nu_L u$	0	$\sqrt{2}g_L$	$g_L < 0.09$	—	—
		$-1/3$	$\nu_L \bar{d}$	$\frac{1}{2}$	$-g_L$		0.40	—
			$e_L u$		$-g_L$			LL
		$-4/3$	$e_L d$	1	$-\sqrt{2}g_L$		0.21	LL
$V_{1/2}$		$-1/3$	$\nu_L \bar{d}$	0	g_L	$g_L < 0.09$	—	—
			$e_R u$	1	g_R	0.30	RL	
		$-4/3$	$e_L \bar{d}$	1	g_L	$g_R < 0.05$	0.32	LR
$e_R d$			g_R	0.32	RL			
$\tilde{V}_{1/2}$		$+2/3$	$\nu_L u$	0	g_L	$g_L < 0.09$	—	—
		$-1/3$	$e_L u$	1	g_L		0.32	LR
$S_{1/2}$		$-2/3$	$\nu_L \bar{u}$	0	g_L	$g_L < 0.1$	—	—
			$e_R \bar{d}$	1	$-g_R$	$g_R < 0.09$	0.052	RL
		$-5/3$	$e_L \bar{u}$	1	g_L		0.026	LR
$e_R \bar{u}$			g_R	0.026	RL			
$\tilde{S}_{1/2}$ *	$\bar{\tilde{d}}_L$	$+1/3$	$\nu_L \bar{d}$	0	g_L	$g_L < 0.1$	—	—
	$\bar{\tilde{u}}_L$	$-2/3$	$e_L \bar{d}$	1	g_L		0.052	LR
V_0		$-2/3$	$e_L \bar{d}$	$\frac{1}{2}$	g_L	$g_L < 0.05$	0.080	LL
			$\nu_L \bar{u}$		g_L	$g_R < 0.09$	0.056	—
			$e_R \bar{d}$	1	g_R			RR
$\tilde{V}_0$		$-5/3$	$e_R \bar{u}$	1	g_R	$g_R < 0.09$	0.027	RR
V_1		$+1/3$	$\nu_L \bar{d}$	0	$\sqrt{2}g_L$	$g_L < 0.04$	—	—
		$-2/3$	$e_L \bar{d}$	$\frac{1}{2}$	$-g_L$		0.080	LL
			$\nu_L \bar{u}$		g_L			—
		$-5/3$	$e_L \bar{u}$	1	$\sqrt{2}g_L$		0.019	LL

Table 1: Scalar (S) and vector (V) leptoquarks/squarks with electric charges (Q), decay modes, branching ratios for charged lepton + jet channels with either L or R couplings, and the Yukawa couplings ($g_{R,L}$) with the most stringent limits from rare decays and estimates from the recent HERA data (see text). The helicity combinations ik ($= L, R$) contributing to the process $e_i^- e^+ \rightarrow q_k \bar{q}$ are given in the last column. Also shown are possible squark assignments of leptoquark-type states; the special leptoquark singled out in Ref. [7] is marked by an asterisk.

variety of phenomena are expected to be observed experimentally. In electron/positron–proton collisions, these particles are produced as single resonances, with the rate determined by the strength of the LQ - l - q Yukawa couplings [8, 14]. Pair production is an important production mechanism for leptoquarks in proton–(anti)proton [15], electron–positron [16] and photon–photon collisions [17]. In these reactions, the size of the cross section is determined (*modulo* anomalous couplings and form-factor effects) by the color, the electric and the electroweak charges of a given leptoquark (if the Yukawa couplings are small [18]). The predictions for the cross sections are therefore, to a first approximation, model-independent. Associated production of $LQ+l$ or $LQ+q$, again mediated by Yukawa interactions, can also be explored in these and other collision processes ($e\gamma$, for example [19]).

In addition to the direct production of leptoquarks/squarks, which is of course of central interest, indirect effects generated by the exchange of virtual LQ s, are also important experimental tools to provide cross checks, to explore the nature of these particles, and to have a glimpse at states which are too heavy to be produced directly. Virtual leptoquarks could strongly affect decay processes such as $\pi \rightarrow l\nu$ and $K \rightarrow \pi\nu\bar{\nu}$. These processes are suppressed for the LQ s considered here by the restrictions on the Yukawa couplings summarized above. Nevertheless, they still contribute to rare pion and kaon decays, $K^0 - \bar{K}^0$ and $D^0 - \bar{D}^0$ mixing, and atomic parity violation [10, 11, 12]. The strongest constraints can be deduced from atomic parity violation with two exceptions: the constraints on $g_L(S_0)$ and $g_L(V_0)$ which follow from the violation of universality in $\pi \rightarrow l\nu$ [11] or in μ and β decays [12, 13]. The bounds, as derived in Refs. [11] and [12], are presented in Table 1 for the reference mass $m_{LQ} = 200$ GeV; the leptoquarks are taken mass degenerate within the isospin multiplets. In the few hundred GeV range the bounds on the couplings scale linearly with the leptoquark mass. Potential destructive interference effects between different states can lift these bounds considerably.

Additional indirect constraints can be derived from the t/u channel exchange of leptoquarks/squarks in high energy processes, such as Drell–Yan production of lepton pairs in pp and $p\bar{p}$ collisions [20], or e^+e^- annihilation to hadrons [21]. The existing bounds can be improved significantly with the rise of the e^+e^- energy to the LEP2 value close to $\sqrt{s} = 200$ GeV.

In the present analysis we study the production of $q\bar{q}$ pairs in e^+e^- annihilation,

$$e^+e^- \rightarrow q\bar{q} \tag{1}$$

which is mediated by γ, Z exchanges in the s -channel (Fig. 1a), and leptoquark/squark exchanges in the t/u -channels (Fig. 1b and c). By definition, t -channel exchange is associated with $e^- \rightarrow q$ transitions ($F = 0$), while u -channel exchange involves $e^- \rightarrow \bar{q}$ transitions ($F = 2$). We will consider both scalar and vector leptoquarks. Earlier analyses of Ref. [21] will be extended in several aspects: (i) We present a systematic analysis of the exchange of all types of leptoquarks. In particular, generalizing the helicity method of Ref. [22], a representation of the exchange amplitudes can be derived in which potential interference effects between different types of leptoquarks are made transparent. (ii) Simple expressions




Figure 1: *Feynman diagrams for $e^+e^- \rightarrow q\bar{q}$ including leptoquarks and squarks in R -parity breaking supersymmetric theories.*

for the integrated cross sections are presented. (iii) Special emphasis is given to squark exchange mechanisms in supersymmetric theories with R -parity breaking, complementing earlier discussions in Ref. [23].

2 Leptoquark/Squark Exchange in e^+e^- Annihilation

After performing the standard Fierz transformations from t/u -channel leptoquark/squark exchange amplitudes in $e^+e^- \rightarrow q\bar{q}$ to standard s -channel amplitudes, only terms of the structure *(lepton vector current) $\times$ (quark vector current)* are generated for leptoquarks carrying either left *or* right chiral couplings, but not *both* at the same time. This leads to a transparent representation of the matrix elements. Denoting the helicity amplitudes of the process $e_i^- e^+ \rightarrow q_k \bar{q}$ with $i, k = R, L$ by f_{ik} and the spin-density matrix elements, which depend on the polar angle θ between e^- and q , by ρ_{ik} , the cross section for the process Eq. (1) can be written as

$$\frac{d\sigma}{d\cos\theta}(e^+e^- \rightarrow q\bar{q}) = \frac{N_c}{128\pi s} \sum_{i,k=L,R} \rho_{ik} |f_{ik}|^2 \quad (2)$$

with the color factor $N_c = 3$. As a consequence of angular momentum conservation, the spin-density matrix elements are given by

$$\rho_{RR} = \rho_{LL} = s^2(1 + \cos\theta)^2 \quad (3)$$

$$\rho_{RL} = \rho_{LR} = s^2(1 - \cos\theta)^2 \quad (4)$$

The helicity amplitudes can be cast in the following form

$$f_{RR} = \frac{Q_{RR}^{eq}}{s} + \delta_{qu} \left(\frac{g_R^2}{t - \tilde{V}_0} - \frac{g_R^2}{2(u - S_0)} \right) + \delta_{qd} \left(\frac{g_R^2}{t - V_0} - \frac{g_R^2}{2(u - \tilde{S}_0)} \right)$$

$$\begin{aligned}
f_{LL} &= \frac{Q_{LL}^{eq}}{s} + \delta_{qu} \left(\frac{2g_L^2}{t-V_1} - \frac{g_L^2}{2(u-S_0)} - \frac{g_L^2}{2(u-S_1)} \right) + \delta_{qd} \left(\frac{g_L^2}{t-V_0} + \frac{g_L^2}{t-V_1} - \frac{g_L^2}{u-S_1} \right) \\
f_{RL} &= \frac{Q_{RL}^{eq}}{s} + \delta_{qu} \left(\frac{g_R^2}{2(t-S_{1/2})} - \frac{g_R^2}{u-V_{1/2}} \right) + \delta_{qd} \left(\frac{g_R^2}{2(t-S_{1/2})} - \frac{g_R^2}{u-V_{1/2}} \right) \\
f_{LR} &= \frac{Q_{LR}^{eq}}{s} + \delta_{qu} \left(\frac{g_L^2}{2(t-S_{1/2})} - \frac{g_L^2}{u-\tilde{V}_{1/2}} \right) + \delta_{qd} \left(\frac{g_L^2}{2(t-\tilde{S}_{1/2})} - \frac{g_L^2}{u-V_{1/2}} \right)
\end{aligned} \tag{5}$$

To simplify the notation in the above formulae, we have used the LQ symbols to denote the mass squared, $LQ := m_{LQ}^2$; the couplings are denoted generically by g_R or g_L with indices identifying the leptoquark type suppressed. The generalized charges in the standard γ, Z exchange amplitudes have been abbreviated by Q_{ik}^{eq} where

$$Q_{ik}^{eq} = e^2 Q_e Q_q + \frac{g_i^e g_k^q}{1 - m_Z^2/s} \tag{6}$$

with the left/right Z charges of the fermions defined as

$$\begin{aligned}
g_L^f &= \frac{e}{s_W c_W} [I_3^f - s_W^2 Q_f] \\
g_R^f &= \frac{e}{s_W c_W} [-s_W^2 Q_f]
\end{aligned}$$

and $s_W = \sin \Theta_W$, $c_W = \cos \Theta_W$. The Mandelstam variables t, u can be expressed by the production angle θ : $t = -s(1 - \cos \theta)/2$, $u = -s(1 + \cos \theta)/2$; they are both negative so that the amplitudes for LQ exchange do not change the sign when θ is varied from the forward to the backward direction.

It is obvious from the expressions (5) that leptoquarks/squarks of a given fermion number F contribute with a fixed positive or negative sign to the helicity amplitude f_{ik} , thus reinforcing their impact mutually. For a given F , the sign of the interference with γ/Z exchange is determined by the sign of the generalized charges Q_{ik}^{eq} , which, in the energy range considered here, are negative for u -quarks and positive for d -quarks, except Q_{RL}^{eq} which is negative. By contrast, leptoquarks with different fermion numbers interfere destructively with each other. Leptoquarks with integer isospin contribute to equal-helicity LL and RR amplitudes, while leptoquarks with $I = 1/2$ contribute to opposite-helicity amplitudes RL and LR .

The angular integration can easily be performed to obtain the total cross section for $e^+e^- \rightarrow q\bar{q}$ including the exchange of one leptoquark with either left or right coupling:

$$\sigma(e^+e^- \rightarrow q\bar{q}) = \frac{N_c}{128\pi s} \left[\frac{8}{3} (|Q_{RR}^{eq}|^2 + |Q_{LL}^{eq}|^2 + |Q_{RL}^{eq}|^2 + |Q_{LR}^{eq}|^2) + \sum_{i=1}^4 k_i C_i(\tau) \right] \tag{7}$$

The ratio τ is defined as $\tau = m_{LQ}^2/s$. If two or more leptoquarks contribute to the same helicity amplitude, interference terms between pairs of leptoquarks must be included;

LQ	$u\bar{u}$ final state				$d\bar{d}$ final state			
	k_1	k_2	k_3	k_4	k_1	k_2	k_3	k_4
S_0		$-Q_{RR}^{eu}g_R^2$		$\frac{1}{4}g_R^4$				
S_0		$-Q_{LL}^{eu}g_L^2$		$\frac{1}{4}g_L^4$				
$\tilde{S}_0$						$-Q_{RR}^{ed}g_R^2$		$\frac{1}{4}g_R^4$
S_1		$-Q_{LL}^{eu}g_L^2$		$\frac{1}{4}g_L^4$		$-2Q_{LL}^{ed}g_L^2$		g_L^4
$V_{1/2}$					$-2Q_{RL}^{ed}g_R^2$		g_R^4	
$V_{1/2}$	$-2Q_{RL}^{eu}g_R^2$		g_R^4		$-2Q_{LR}^{ed}g_L^2$		g_L^4	
$\tilde{V}_{1/2}$	$-2Q_{LR}^{eu}g_L^2$		g_L^4					
$S_{1/2}$		$Q_{RL}^{eu}g_R^2$		$\frac{1}{4}g_R^4$		$Q_{RL}^{ed}g_R^2$		$\frac{1}{4}g_R^4$
$S_{1/2}$		$Q_{LR}^{eu}g_L^2$		$\frac{1}{4}g_L^4$				
$\tilde{S}_{1/2}$						$Q_{LR}^{ed}g_L^2$		$\frac{1}{4}g_L^4$
V_0					$2Q_{RR}^{ed}g_R^2$		g_R^4	
V_0					$2Q_{LL}^{ed}g_L^2$		g_L^4	
$\tilde{V}_0$	$2Q_{RR}^{eu}g_R^2$		g_R^4					
V_1	$4Q_{LL}^{eu}g_L^2$		$4g_L^4$		$2Q_{LL}^{ed}g_L^2$		g_L^4	

Table 2: The coefficients k_i in Eq. (7) describing the exchange of leptoquarks in the total cross section of e^+e^- annihilation to hadrons.

they are collected in the Appendix. The interference terms between leptoquark and γ, Z exchange amplitudes are described by two functions, depending on the mass ratio τ ,

$$C_1(\tau) = 12 + 8\tau - 8(1 + \tau)^2 \log \frac{1 + \tau}{\tau} \quad (8)$$

$$C_2(\tau) = 8\tau - 4 - 8\tau^2 \log \frac{1 + \tau}{\tau} \quad (9)$$

The couplings building up k_1 and k_2 are listed in Table 2. The squared leptoquark-exchange amplitudes are given by two additional functions,

$$C_3(\tau) = 16 + \frac{8}{\tau} - 16(1 + \tau) \log \frac{1 + \tau}{\tau} \quad (10)$$

$$C_4(\tau) = 16 - \frac{8}{1 + \tau} - 16\tau \log \frac{1 + \tau}{\tau} \quad (11)$$

with the coefficients k_3 and k_4 again listed in Table 2.

It is instructive to consider the helicity amplitudes explicitly in the large mass limit $m_{LQ} \gg \sqrt{s}$. In this limit they can be interpreted as lepton-quark contact terms:

$$\begin{aligned}
f_{RR} &= \frac{Q_{RR}^{eq}}{s} - \delta_{qu} \left(\frac{g_R^2}{\tilde{V}_0} - \frac{g_R^2}{2S_0} \right) - \delta_{qd} \left(\frac{g_R^2}{\tilde{V}_0} - \frac{g_R^2}{2\tilde{S}_0} \right) \\
f_{LL} &= \frac{Q_{LL}^{eq}}{s} - \delta_{qu} \left(\frac{2g_L^2}{V_1} - \frac{g_L^2}{2S_0} - \frac{g_L^2}{2S_1} \right) - \delta_{qd} \left(\frac{g_L^2}{\tilde{V}_0} + \frac{g_L^2}{V_1} - \frac{g_L^2}{S_1} \right) \\
f_{RL} &= \frac{Q_{RL}^{eq}}{s} - \delta_{qu} \left(\frac{g_R^2}{2S_{1/2}} - \frac{g_R^2}{V_{1/2}} \right) - \delta_{qd} \left(\frac{g_R^2}{2S_{1/2}} - \frac{g_R^2}{V_{1/2}} \right) \\
f_{LR} &= \frac{Q_{LR}^{eq}}{s} - \delta_{qu} \left(\frac{g_L^2}{2S_{1/2}} - \frac{g_L^2}{\tilde{V}_{1/2}} \right) - \delta_{qd} \left(\frac{g_L^2}{2\tilde{S}_{1/2}} - \frac{g_L^2}{V_{1/2}} \right)
\end{aligned} \tag{12}$$

As observed before, leptoquarks with integer isospin $I = 0$ and $I = 1$ build up equal-helicity LL and RR contact terms, while leptoquarks with $I = 1/2$ contribute to opposite-helicity RL and LR contact terms. These rules may also be cast into the standard form of the effective Lagrangian [24]

$$\begin{aligned}
\mathcal{L}_{eff} &= \sum_{i,k=L,R} \frac{g_i^2}{m_{LQ}^2} \alpha^{ik} (\bar{e}_i \gamma^\mu e_i) (\bar{q}_k \gamma_\mu q_k) \\
&:= \sum_{i,k=L,R} \eta_{ik} \frac{4\pi}{\Lambda_{ik}^2} (\bar{e}_i \gamma^\mu e_i) (\bar{q}_k \gamma_\mu q_k)
\end{aligned} \tag{13}$$

with e_i, q_k denoting left- and right-handed electron and quark fields. The coefficients α^{ik} for $u\bar{u}$ and $d\bar{d}$ final states are listed in Table 3. Denoting the signs of α^{ik} by η_{ik} , the scales Λ_{ik}^2 of the contact interactions are related to the individual masses and couplings of the leptoquarks by $\Lambda_{ik}^2 = 4\pi m_{LQ}^2 / g_i^2 |\alpha^{ik}|$. In the total cross section $\sigma(e^+e^- \rightarrow q\bar{q})$ the interference terms and the squared contact terms approach the limits $C_1 = C_2 = -8s/3m_{LQ}^2$ and $C_3 = C_4 = 8s^2/3m_{LQ}^4$ for $m_{LQ} \gg \sqrt{s}$.

3 Phenomenological Evaluation

If the surplus of the HERA high Q^2 , large x events is interpreted as the production of scalar or vector leptoquarks, or of squarks, their Yukawa couplings can be estimated from the production rates. We present only *qualitative estimates* of these couplings which should illustrate the general expectations for possible effects in e^+e^- annihilation but which should not anticipate a rigorous analysis to be performed by the experiments themselves. Nevertheless, averaging over the H1 and ZEUS data one finds the couplings listed in Table 1. It is assumed in these estimates that only one type of leptoquark (*i.e.* a single member of an isomultiplet) has been generated with one specific chiral coupling (L, R) to one specific quark flavor (*up, down*) which gives rise to the branching ratios for the decays of leptoquarks into charged leptons shown in the same Table. The $F = 2$ leptoquarks (upper part

α^{ik}	$u\bar{u}$ final state				$d\bar{d}$ final state			
	RR	LL	RL	LR	RR	LL	RL	LR
S_0	$\frac{1}{2}$	$\frac{1}{2}$						
$\tilde{S}_0$					$\frac{1}{2}$			
S_1		$\frac{1}{2}$				1		
$V_{1/2}$			1				1	1
$\tilde{V}_{1/2}$				1				
$S_{1/2}$			$-\frac{1}{2}$	$-\frac{1}{2}$			$-\frac{1}{2}$	
$\tilde{S}_{1/2}$								$-\frac{1}{2}$
V_0					-1	-1		
$\tilde{V}_0$	-1							
V_1		-2				-1		

Table 3: *The coefficients α^{ik} in the Lagrangian of the contact interactions.*

of the Table) are generated in positron sea-quark collisions, the $F = 0$ leptoquarks (lower part of the Table) in positron valence-quark collisions.

From the couplings shown in Table 1 we can draw interesting conclusions even at the present level of qualitative arguments:

(i) If the HERA events are interpreted as the signal of a leptoquark generated in positron valence-quark collisions, the Yukawa coupling is of the order $\sim e/10$, *i.e.* suppressed by a full order of magnitude compared with the electromagnetic coupling. The t/u -channel exchange of such a state affects the $e^+e^- \rightarrow u\bar{u}$ or $e^+e^- \rightarrow d\bar{d}$ parton cross sections generally only at the level of a percent (up to 10 % for V_0 and V_1). If all parton channels are summed up, the impact on the total hadronic cross section is even smaller if the heavier flavors are not excited by the contact interactions.

(ii) The coupling for single leptoquark production out of the sea in positron scattering is large, of order e . However, for such couplings the cross section $\sigma(e^-p \xrightarrow{q} LQ)$ would be two orders of magnitude larger than $\sigma(e^+p \xrightarrow{\bar{q}} LQ)$ and it seems unlikely that the large number of events with $m_{LQ} \sim 200$ GeV could have been missed in the earlier electron-proton runs at HERA². Nevertheless, Yukawa couplings of the order $g_{L,R} \sim e/3$ do not seem to be ruled out yet completely [25]. In such a scenario several types of leptoquarks

²Moreover, the low energy bounds restrict the couplings to $g_{L,R} \lesssim 0.1$, see Table 1.

could be responsible for the observed events at HERA. The $F = 2$ leptoquarks may lead to observable effects in hadron production in e^+e^- annihilation at LEP2, or at least more stringent bounds on the Yukawa couplings can be established.

(iii) If the HERA events are interpreted as the signal of leptoquark resonance production, they must originate from valence quarks, that is from e^+u or e^+d fusion to $F = 0$ states, and they must couple chirally to an extremely good approximation in order to be consistent with the existing bounds. Because of charge conservation, there is only one possible process which can give rise to new events in charged-current reactions, $e^+d \rightarrow \bar{\nu}u$, and only in the presence of left-handed couplings. This implies that an excess of events in the CC channel, in addition to the NC channel, will single out just two states: the vector LQ s V_0 and V_1 carrying charge $\frac{2}{3}$. Thus, even without switching from an e^+ to an e^- beam, the NC and CC searches combined are very selective in the LQ quantum numbers.

However, recalling that no symmetry principles are known which give rise to relations among Yukawa scalar-fermion couplings – even within isomultiplets they differ by nearly two orders of magnitude in the Higgs sector – there could be additional leptoquarks at higher masses and with larger couplings that cannot be observed as resonance states in the HERA experiments. We have therefore studied the sensitivity of the total hadronic e^+e^- cross section to the entire ensemble of scalar and vector leptoquarks listed in Table 1. The result³ is illustrated in Figs. 2 and 3 for the parameter set $(g_L, g_R) = (0.1, 0)$ or $(0, 0.1)$ at $\sqrt{s} = 192$ GeV as a function of the LQ mass. Since the couplings are arbitrary parameters, only the relative size of the curves is relevant. For small enough couplings and large enough masses the curves scale in $g_{L,R}^2/m_{LQ}^2$. We observe both constructive and destructive interference effects, depending on the type of quarks in the final state. The impact of $I = 0, 1$ leptoquarks on the hadronic cross section is larger than the impact of $I = 1/2$ leptoquarks.

For large masses, the exchange of leptoquarks can be described by contact interactions. Depending on the type of leptoquark, different helicity combinations of lepton and quark currents are affected in either $u\bar{u}$ or $d\bar{d}$ final states. The sign of the contact interactions depends on the fermion number as shown in the previous section. Potentially large effects can be expected for e^+e^- annihilation to hadrons. This is exemplified for a series of Λ values in Table 4. The symbols LL etc. denote the helicity of the lepton current followed by the helicity of the quark current. The effect of the contact interaction (CI) is shown for the total e^+e^- hadronic cross section at LEP192 if only one of the u -type or d -type quarks is involved in the contact interactions. Present analyses of hadron production at LEP set limits to Λ already at the level of about 1.5 to 2.5 TeV [27].

³Numerical cross checks have been performed with the help of CompHEP [26] adapted to leptoquark processes.




Figure 2: *Effect of t/u -channel exchange of scalar leptoquarks on the total hadronic cross section as a function of m_{LQ} for $\sqrt{s} = 192$ GeV. The couplings have been fixed arbitrarily to $(g_L, g_R) = (0.1, 0)$ or $(0, 0.1)$ indicated by $LQ^{L,R}$, respectively.*




Figure 3: *Effect of t/u -channel exchange of vector leptoquarks on the total hadronic cross section as a function of m_{LQ} for $\sqrt{s} = 192$ GeV. The couplings have been fixed arbitrarily to $(g_L, g_R) = (0.1, 0)$ or $(0, 0.1)$ indicated by $LQ^{L,R}$, respectively.*

ik $\Lambda[\text{GeV}]$	$u\bar{u}$ final state			$d\bar{d}$ final state		
	1.5	2.5	3.5	1.5	2.5	3.5
	$\eta = +1$					
LL	-0.11	-0.14	-0.09	0.92	0.23	0.10
RR	0.08	-0.07	-0.05	0.63	0.12	0.05
LR	0.30	0.01	-0.01	0.52	0.08	0.003
RL	0.41	0.05	0.01	0.30	0.004	-0.001
	$\eta = -1$					
LL	0.99	0.26	0.12	-0.03	-0.11	-0.07
RR	0.81	0.19	0.08	0.26	-0.008	-0.02
LR	0.59	0.11	0.04	0.37	0.03	0.002
RL	0.48	0.07	0.02	0.59	0.11	0.04

Table 4: The effect of contact interactions with different helicities on the cross section for hadron production in e^+e^- annihilation: $\Delta = \sigma(SM \oplus CI)/\sigma(SM) - 1$.

4 Squarks in R -parity Breaking SUSY Models

In the minimal supersymmetric extension of the Standard Model, the only renormalizable, gauge invariant operator that couples squarks to quarks and leptons is given by

$$W_R = \lambda'_{ijk} L_L^i Q_L^j \bar{D}_R^k \quad (14)$$

in the superpotential [6]. The indices ijk are generation indices in the left-handed doublets of leptons (L) and quarks (Q), and right-handed singlets of down-type quarks (D). This interaction term violates global invariance of R -parity, defined as $R = (-1)^{3B+L+2S}$ which is $+1$ for particles and -1 for superpartners. This interaction has also been considered in the context of the Aleph 4-jet events in Ref. [29].

Expanding the superfields in terms of matter fields, the interaction Lagrangian can be written as

$$\mathcal{L}_R = \lambda'_{ijk} \left[\tilde{u}_L^j \bar{d}_R^k e_L^i + \bar{\tilde{d}}_R^k (\bar{e}_L^i)^c u_L^j + \tilde{e}_L^i \bar{d}_L^k u_L^j - \tilde{\nu}_L^i \bar{d}_R^k d_L^j - \tilde{d}_L^j \bar{d}_R^k \nu_L^i - \bar{\tilde{d}}_R^k (\bar{\nu}_L^i)^c d_L^j \right] + h.c. \quad (15)$$

where u^j and d^k stand for u - and d -type quarks, respectively, and the superscript $(\)^c$ denotes charge conjugate spinors. The first two terms with $i = 1$ are of particular interest since they allow for resonant squark production in e^+p scattering at HERA via the subprocesses

$$e^+ + d_R^k \rightarrow \tilde{u}_L^j \quad (\tilde{u}^j = \tilde{u}, \tilde{c}, \tilde{t}) \quad (16)$$

$$e^+ + \bar{u}_L^j \rightarrow \bar{\tilde{d}}_R^k \quad (\tilde{d}^k = \tilde{d}, \tilde{s}, \tilde{b}) \quad (17)$$

If more than one of the couplings λ'_{1jk} is non-vanishing, strong limits on their product are




Figure 4: *Squarks production mechanisms in e^+p collisions.*

imposed by the absence of FCNC reactions [28]. Since $\lambda'_{111} \lesssim 10^{-3}$ (for a mass $m = 200$ GeV of supersymmetric partners mediating neutrinoless double beta decay [13]), second or third generation fermions must be coupled to electrons in order to account for the rate at HERA. Below we will consider the two possible scenarios, shown in Fig. 4, in which a) quarks, b) antiquarks from the proton participate in the production process. In particular we will discuss two cases for $\lambda'_{1j1} \neq 0$ or $\lambda'_{11k} \neq 0$ and their implications for e^+e^- annihilation to hadrons.

a) $\lambda'_{1j1} \neq 0, j = 2, 3$




Figure 5: The scenario with $\lambda'_{1j1} \neq 0$: $\tilde{c}$ for $j = 2$, $\tilde{t}$ for $j = 3$.

In this case, *down* quarks are involved⁴ via (16) in the production of left-handed *charm* ($j = 2$) or *top* ($j = 3$) squarks at HERA, Fig. 4a. The other process (17) is irrelevant since

⁴Another possibility is to consider $\lambda'_{132} \neq 0$ which would involve *strange* quarks from the sea in the production of *top* squarks (Fig. 4a). The qualitative analysis below applies to this case as well: $s\bar{s}$ production via $\tilde{t}_L$ (see Fig. 5a and Fig. 6).

it would require *charm* or *top* sea antiquarks in the proton. To account for the observed number of events at HERA, λ_{1j1} must exceed 0.052 which is still within the limits derived from atomic parity violation for a sufficiently heavy $\tilde{d}_R$ [11, 28]. Notice that since $\tilde{u}^j$ does not couple to neutrinos, the surplus of events in CC reactions at HERA is not expected. In $e^+e^- \rightarrow q\bar{q}$ the coupling λ'_{1j1} leads to two additional hadron channels, as shown in Fig. 5: $e^+e^- \rightarrow d\bar{d}$ via *charm* or *top* squark exchange in the t -channel, and $e^+e^- \rightarrow c\bar{c}$ via *down* squark exchange in the u -channel. Since the left-handed *up*-type squarks couple in the same way as the $\tilde{S}_{1/2}$ leptoquark, and the right-handed *down* squark like the S_0 leptoquark with left-handed coupling g_L , their contributions can easily be obtained from the formulae given in the previous section. The impact of these exchange mechanisms on the total hadronic e^+e^- cross section is shown in Fig. 6. The impact of squark exchange on the single parton cross section $e^+e^- \rightarrow c\bar{c}$ is larger; the experimental analysis of this process however requires the tagging of *charm* quarks in the final state.

b) $\lambda'_{11k} \neq 0, k = 2, 3$

In this case, *up* antiquarks of the sea are involved in the process (17) so that *strange* ($k = 2$) or *bottom* ($k = 3$) antisquarks would be produced at HERA, Fig. 4b. The $\tilde{d}$ couples also to neutrinos, therefore similar events in CC reactions could be expected. The coupling λ'_{11k} would introduce two additional mechanisms in the $e^+e^- \rightarrow q\bar{q}$ process, Fig. 7: $e^+e^- \rightarrow s\bar{s}$ or $e^+e^- \rightarrow b\bar{b}$ via *up* squark exchange in the t -channel, and $e^+e^- \rightarrow u\bar{u}$ via *strange* or *bottom* squark exchange in the u -channel. Since the parton densities of sea quarks are much smaller than the densities of valence quarks, this scenario would require a large coupling, $\lambda'_{11k} > 0.3$. Such a large value of λ'_{11k} is however in conflict with the limit $\lambda'_{11k} < 0.06$ derived from charged current universality [30] or from earlier e^-p data [25]. This mechanism therefore cannot explain the surplus of HERA data.

In contrast to genuine leptoquarks which decay solely to leptons and quarks, squarks can decay not only via R -parity violating couplings but in general also via a large number of R -parity conserving modes: $\tilde{q} \rightarrow q\chi$ with χ being either a neutralino or a chargino state, cascading in a chain reaction down to ordinary particles. The lower limits for the couplings λ' inferred from the HERA events were based on branching ratios of 100 % for the R -parity violating decay modes to lepton plus quark jet. If the branching ratios for $\tilde{q} \rightarrow q\chi$ decays are non-negligible, the couplings λ' would have to be larger correspondingly. This would increase the impact on e^+e^- collisions. At the same time, new types of events at HERA would be expected with multiple jet topologies and leptons from the R -parity breaking χ decays.




Figure 6: *Effect of t/u -channel exchange of squarks in the supersymmetry scenario (a) on the total hadronic cross section, $\Delta = \sigma(SM \oplus \tilde{q})/\sigma(SM) - 1$, as a function of $m_{\tilde{q}}$ for $\lambda'_{1j1} = 0.1$, $j = 2$ or 3 (or $\lambda'_{132} = 0.1$ for $s\bar{s}$) and $\sqrt{s} = 192$ GeV.*




Figure 7: *The scenario with $\lambda'_{11k} \neq 0$: $\tilde{s}$ for $k = 2$, $\tilde{b}$ for $k = 3$.*

5 Summary

The conclusions of our analysis can be summarized in four points.

(i) If the high Q^2 , large x events observed at HERA in deep-inelastic positron-proton scattering are interpreted as the direct production of a narrow leptoquark state, two cases must be distinguished. For leptoquarks generated in collisions with valence quarks, the Yukawa couplings are so small, $\sim e/10$, that the contribution of the t/u -channel exchange of these leptoquarks affects the production of hadrons in high-energy e^+e^- annihilation only at the level of less than one percent. For leptoquarks generated from antiquarks in the sea, the Yukawa couplings can still be larger. However, the couplings presumably do not exceed the value $e/3$ since the leptoquark states would have been observed otherwise in the earlier electron-proton runs at HERA. In this second case, the effects of leptoquark exchange may be accessible in e^+e^- annihilation at LEP2.

(ii) For masses above the range covered directly by HERA, the impact of the leptoquark exchange on hadron production in e^+e^- annihilation can be significant for a wide range of Yukawa couplings. The interactions are effectively described by contact terms, similar to contact interactions at HERA. The cross sections have been presented for all standard leptoquark states.

(iii) In R -parity breaking supersymmetric models, the HERA events could be interpreted as the production of either left-handed *charm* or *top* squarks. The observed number of events requires $\lambda'_{1j1} > 0.065$, the lower limit corresponding to a branching ratio of $\sim 100\%$ for the R -parity violating decays to lepton and hadron jets. If the coupling is close to the lower limit, the impact on the hadronic cross section in e^+e^- annihilation is small, $\sim 1\%$. If the coupling is stronger, the impact on e^+e^- annihilation will be more pronounced. In this case, interesting multi-jet and lepton signatures due to the R -parity conserving decays $\tilde{q} \rightarrow q\chi$, followed by R -parity breaking χ decays, could occur at HERA.

(iv) If the HERA events are not interpreted as the production of narrow leptoquark reso-

nances but are described globally by contact interactions, the effective scale Λ is predicted to be in a range which is easily accessible at LEP2. However, if deviations from the prediction of the Standard Model for the cross section of e^+e^- annihilation to hadrons are not observed, contact interactions with scales of order 2 TeV cannot account for the HERA events. This would restrict the interpretation of the events to the direct production of narrow leptoquark-type resonances.

Appendix

If more than one leptoquark contributes to the same helicity amplitude, the expression in Eq. (7) has to be supplemented by the interference terms between pairs of leptoquarks. For the $u\bar{u}$ final states we find

$$\frac{N_c}{128\pi s} \left[+\frac{1}{2}g_{1L}^2g_{3L}^2C_6(\tau_1, \tau_3) - g_{1R}^2g_{9R}^2C_7(\tau_9, \tau_1) - 2g_{10L}^2g_{1L}^2C_7(\tau_{10}, \tau_1) \right. \quad (18)$$

$$\left. -2g_{10L}^2g_{3L}^2C_7(\tau_{10}, \tau_3) - g_{6R}^2g_{4R}^2C_7(\tau_4, \tau_6) - g_{6L}^2g_{5L}^2C_7(\tau_5, \tau_6) \right] \quad (19)$$

and for $d\bar{d}$ final states

$$\frac{N_c}{128\pi s} \left[+2g_{8L}^2g_{10L}^2C_5(\tau_{10}, \tau_8) - g_{2R}^2g_{8R}^2C_7(\tau_8, \tau_2) - 2g_{8L}^2g_{3L}^2C_7(\tau_8, \tau_3) \right. \quad (20)$$

$$\left. -2g_{10L}^2g_{3L}^2C_7(\tau_{10}, \tau_3) - g_{6R}^2g_{4R}^2C_7(\tau_4, \tau_6) - g_{7L}^2g_{4L}^2C_7(\tau_4, \tau_7) \right] \quad (21)$$

The numbering of the leptoquark states and their couplings (g_{iL} or g_{iR}) and masses ($\tau_i = m_i^2/s$) follows the listing in Table 1, with $i = 1, \dots, 10$. For example g_{6R} and τ_6 refer to the right-handed coupling and mass of the $S_{1/2}$ state. The functions C_5 , C_6 and C_7 can be expressed in terms of C_1 and C_2 as follows:

$$\begin{aligned} C_5(x, y) &= \frac{C_1(x) - C_1(y)}{2y - 2x} \\ C_6(x, y) &= \frac{C_2(x) - C_2(y)}{2y - 2x} \\ C_7(x, y) &= \frac{C_1(x) + C_2(y)}{2 + 2x + 2y} \end{aligned} \quad (22)$$

Acknowledgments:

We have benefitted from discussions with E. Boos, W. Buchmüller, P. Mättig and U. Martyn.

NOTE: When finalizing the present paper we received copies of Refs. [31, 32]. The main conclusions are in mutual agreement.

References

- [1] H1 Collab., C. Adloff et al., DESY 97-024 and *Z. Phys. C*, in press;
ZEUS Collab., J. Breitweg et al., DESY 97-025 and *Z. Phys. C*, in press.
- [2] J.C. Pati and A. Salam, *Phys. Rev.* **D8** (1973) 1240, *ibid.* **10** (1974) 275.
- [3] H. Georgi and S.L. Glashow, *Phys. Rev. Lett.* **32** (1974) 438; P. Langacker, *Phys. Rep.* **72** (1981) 185; J. L. Hewett and T. G. Rizzo, *Phys. Rep.* **183** (1989) 193.
- [4] L. Abbott and E. Farhi, *Phys. Lett.* **B101** (1981) 69, *Nucl. Phys.* **B189** (1981) 547; W. Buchmüller, *Acta Phys. Austr. Suppl.* XXVII (1985) 517; B. Schrempp and F. Schrempp, *Phys. Lett.* **B153** (1985) 101.
- [5] P. Fayet, *Phys. Lett.* **B69** (1977) 489; G. Farrar and P. Fayet, *Phys. Lett.* **B76** (1978) 575; N. Sakai and T. Yanagida, *Nucl. Phys.* **B197** (1982) 533.
- [6] C.S. Aulakh and R.N. Mohapatra, *Phys. Lett.* **B119** (1982) 316; F. Zwirner, *Phys. Lett.* **B132** (1983) 103; L.J. Hall and M. Suzuki, *Nucl. Phys.* **B231** (1984) 419; S. Dawson, *Nucl. Phys.* **B261** (1985) 297.
- [7] H. Murayama and T. Yanagida, *Mod. Phys. Lett.* **A7** (1992) 147.
- [8] W. Buchmüller, R. Rückl and D. Wyler, *Phys. Lett.* **B191** (1987) 442.
- [9] M. Spira, T. Köhler, Diploma Theses, RWTH Aachen (1989); A. Djouadi, T. Köhler, M. Spira and J. Tutas, *Z. Phys.* **C46** (1990) 679; B. Schrempp, *Proceedings, Physics at HERA* (Hamburg 1991), eds. W. Buchmüller and G. Ingelman.
- [10] O. Shanker, *Nucl. Phys.* **B204** (1982) 253; W. Buchmüller and D. Wyler, *Phys. Lett.* **B177** (1986) 377.
- [11] M. Leurer, *Phys. Rev.* **D49** (1994) 333; *ibid.* **50** (1994) 536.
- [12] S. Davidson, D. Bailey and B. Campbell, *Z. Phys.* **C61** (1994) 613.
- [13] M. Hirsch, H.V. Klapdor-Kleingrothaus and S.G. Kovalenko, *Phys. Rev.* **D53** (1996) 1329.
- [14] J.L. Hewett, *Proceedings, 1990 Summer Study on High Energy Physics – Research Directions for the Decade* (Snowmass 1990), ed. E. Berger (World Scientific, 1992); J. Butterworth and H. Dreiner, *Nucl. Phys.* **B397** (1993) 3.
- [15] J.L. Hewett and S. Pakvasa, *Phys. Rev.* **D37** (1988) 3165; O.J.P. Éboli and A.V. Olinto, *Phys. Rev.* **D38** (1988) 3461; A. Dobado, M.J. Herrero and C. Munoz, *Phys. Lett.* **B207** (1988) 97; M. de Montigny and L. Marleau, *Phys. Rev.* **D41** (1990) 3523; B. Dion, L. Marleau and G. Simon, LAVAL-PHY-96-16, hep-ph/9610397;

- T.G. Rizzo, SLAC-PUB-7284, hep-ph/9609267; J. Blümlein, E. Boos and A. Kryukov, DESY 96-174, hep-ph/9610408; B. Dion, L. Marleau, G. Simon and M. de Montigny, LAVAL-PHY-96-17, ALBERTA-THY-39-96, hep-ph/9701285.
- [16] D. Schaile and P.M. Zerwas, Proceedings, *Physics at Future Accelerators* (La Thuile, Geneva 1987), ed. J.H. Mulvey, CERN 87-07; N.D. Tracas and S.D.P. Vlassopoulos, *Phys. Lett.* **B220** (1989) 285; A. Djouadi, M. Spira and P.M. Zerwas, Proceedings, *e^+e^- Collisions at 500 GeV: The Physics Potential* (Munich, Annecy, Hamburg 1991), ed. P. M. Zerwas, DESY 92-123B; J. Blümlein and R. Rückl, *ibid.*; J.E. Cieza Montalvo and O.J.P. Éboli, *Phys. Rev.* **D47** (1993) 837; J. Blümlein and R. Rückl, *Phys. Lett.* **B304** (1993) 337; D. Choudhury, *Phys. Lett.* **B346** (1995) 291; J. Blümlein, E. Boos and A. Kryukov, DESY 96-219, hep-ph/9610506; O.J.P. Éboli, M.C. Gonzalez-Garcia and J.K. Mizukoshi, FTUV/96-82, IFIC/96-91, IFUSP-P 1251, hep-ph/9612254; M.A. Doncheski and S. Godfrey, hep-ph/9612385.
 - [17] J.E. Cieza Montalvo and O.J.P. Éboli, see Ref. [16].
 - [18] J. Blümlein and R. Rückl, see Ref. [16].
 - [19] J.L. Hewett and S. Pakvasa, *Phys. Lett.* **B227** (1989) 178; O.J.P. Éboli et al., *Phys. Lett.* **B311** (1993) 147. N. Nadeau and D. London, *Phys. Rev.* **D47** (1993) 3742; G. Belanger, N. Nadeau and D. London, *Phys. Rev.* **D49** (1994) 3140; M.A. Doncheski and S. Godfrey, *Phys. Rev.* **D49** (1994) 6220, *ibid.* **51** (1995) 1040.
 - [20] R. Rückl and P.M. Zerwas, Proceedings, *Physics at Future Accelerators* (La Thuile, Geneva 1987), ed. J.H. Mulvey, CERN 87-07; G. Bhattacharyya, D. Choudhury and K. Sridhar, *Phys. Lett.* **B349** (1995) 118.
 - [21] J.L. Hewett and T.G. Rizzo, *Phys. Rev.* **D36** (1987) 3367; H. Dreiner et al., *Mod. Phys. Lett.* **A3** (1988) 443; J.E. Cieza Montalvo and O.J.P. Éboli, *Phys. Rev.* **D47** (1993) 837; D. Choudhury, *Phys. Lett.* **B376** (1996) 201; M.S. Berger, IUHET-343, hep-ph/9609517.
 - [22] L.M. Sehgal and P.M. Zerwas, *Nucl. Phys.* **B183** (1981) 417.
 - [23] D. Choudhury and S. Raychaudhuri, CERN-TH/97-26, hep-ph/9702392.
 - [24] E. Eichten, K. Lane and M. Peskin, *Phys. Rev. Lett.* **50** (1983) 811; R. Rückl, *Phys. Lett.* **B129** (1993) 363, *Nucl. Phys.* **B234** (1984) 91; see also the workshop report P. Haberl, F. Schrempf and H.-U. Martyn, in the Proceedings, *Physics at HERA* (Hamburg, 1991), eds. W. Buchmüller and G. Ingelman.
 - [25] H1 Collab., S. Aid et al., *Phys. Lett.* **B369** (1996) 173; ZEUS Collab., M. Derrick et al., *Phys. Lett.* **B306** (1993) 173.
 - [26] P.A. Baikov et al., Proc. X Int. Workshop QFTHEP'95, hep-ph/9701412.

- [27] P. Mättig, Proceedings, *International Conference on High Energy Physics* (Warsaw 1996); OPAL physics note PN 280 and P. Mättig, private communication.
- [28] K. Agashe and M. Graesser, *Phys. Rev.* **D54** (1995) 4445; D. Choudhury and P. Roy, *Phys. Lett.* **B378** (1996) 153.
- [29] M. Carena, G.F. Giudice, S. Lola and C.E.M. Wagner, CERN-TH/96-352, hep-ph/9612334.
- [30] V. Barger, G.F. Giudice and T. Han, *Phys. Rev.* **D40** (1989) 2987.
- [31] G. Altarelli, J. Ellis, G.F. Giudice, S. Lola and M.L. Mangano, CERN-TH/97-40, hep-ph/9703276.
- [32] H. Dreiner and P. Morawitz, hep-ph/9703279.